

# Partition regularity of nonlinear Diophantine equations

Lorenzo Luperi Baglini

University of Vienna

RaTLoCC 2018: Ramsey Theory in Logic, Combinatorics and Complexity - Bertinoro

# Partition regularity of polynomials/equations

In this talk, we will repeatedly talk about the following property:

# Partition regularity of polynomials/equations

In this talk, we will repeatedly talk about the following property:

## Definition

*Let  $P(x_1, \dots, x_n) \in \mathbb{Z}[x_1, \dots, x_n]$ .*

# Partition regularity of polynomials/equations

In this talk, we will repeatedly talk about the following property:

## Definition

Let  $P(x_1, \dots, x_n) \in \mathbb{Z}[x_1, \dots, x_n]$ . We say that the equation  $P(x_1, \dots, x_n) = 0$  is *(weakly) partition regular (PR) on  $\mathbb{N}$*  if

# Partition regularity of polynomials/equations

In this talk, we will repeatedly talk about the following property:

## Definition

Let  $P(x_1, \dots, x_n) \in \mathbb{Z}[x_1, \dots, x_n]$ . We say that the equation  $P(x_1, \dots, x_n) = 0$  is *(weakly) partition regular (PR) on  $\mathbb{N}$*  if it has a monochromatic solution in every finite coloring of  $\mathbb{N}$ , i.e.

$\forall k \in \mathbb{N}, \forall \mathbb{N} = A_1 \cup \dots \cup A_k \exists i \leq k \exists x_1, \dots, x_n \in A_i$  s.t.

$P(x_1, \dots, x_n) = 0$ .

# Partition regularity of polynomials/equations

In this talk, we will repeatedly talk about the following property:

## Definition

Let  $P(x_1, \dots, x_n) \in \mathbb{Z}[x_1, \dots, x_n]$ . We say that the equation  $P(x_1, \dots, x_n) = 0$  is *(weakly) partition regular (PR) on  $\mathbb{N}$*  if it has a monochromatic solution in every finite coloring of  $\mathbb{N}$ , i.e.

$\forall k \in \mathbb{N}, \forall \mathbb{N} = A_1 \cup \dots \cup A_k \exists i \leq k \exists x_1, \dots, x_n \in A_i$  s.t.  
 $P(x_1, \dots, x_n) = 0$ .

## Example

Trivially, for every  $n \in \mathbb{N}$ , the polynomial  $x - n$  is PR.

# Rado's theorem

## Theorem (Schur)

*The polynomial  $x + y - z$  is PR.*

# Rado's theorem

## Theorem (Schur)

*The polynomial  $x + y - z$  is PR.*

## Theorem (Rado)

*A linear Diophantine equation with no constant term*

$$c_1x_1 + \cdots + c_nx_n = 0$$

*is PR on  $\mathbb{N}$  if and only if the following condition is satisfied:*



# Rado's theorem

## Theorem (Schur)

*The polynomial  $x + y - z$  is PR.*

## Theorem (Rado)

*A linear Diophantine equation with no constant term*

$$c_1x_1 + \cdots + c_nx_n = 0$$

*is PR on  $\mathbb{N}$  if and only if the following condition is satisfied:*

- *“There exists a nonempty set  $J \subseteq \{1, \dots, n\}$  such that  $\sum_{j \in J} c_j = 0$ .”*

## Nonlinear results/1

### Theorem (Multiplicative Rado)

*A nonlinear Diophantine equation  $\prod_{i=1}^n x_i^{c_i} = 1$  is PR on  $\mathbb{N}$  if and only if the following condition is satisfied:*

# Nonlinear results/1

## Theorem (Multiplicative Rado)

A nonlinear Diophantine equation  $\prod_{i=1}^n x_i^{c_i} = 1$  is PR on  $\mathbb{N}$  if and only if the following condition is satisfied:

- “There exists a nonempty set  $J \subseteq \{1, \dots, n\}$  such that  $\sum_{j \in J} c_j = 0$ .”

# Nonlinear results/1

## Theorem (Multiplicative Rado)

A nonlinear Diophantine equation  $\prod_{i=1}^n x_i^{c_i} = 1$  is PR on  $\mathbb{N}$  if and only if the following condition is satisfied:

- “There exists a nonempty set  $J \subseteq \{1, \dots, n\}$  such that  $\sum_{j \in J} c_j = 0$ .”

Idea:  $n \rightarrow 2^n$ .

## Nonlinear results/1

### Theorem (Multiplicative Rado)

A nonlinear Diophantine equation  $\prod_{i=1}^n x_i^{c_i} = 1$  is PR on  $\mathbb{N}$  if and only if the following condition is satisfied:

- “There exists a nonempty set  $J \subseteq \{1, \dots, n\}$  such that  $\sum_{j \in J} c_j = 0$ .”

Idea:  $n \rightarrow 2^n$ .

### Theorem (Lefmann)

Let  $k \in \mathbb{N}$ . A Diophantine equation of the form  $c_1 x_1^{1/k} + \dots + c_n x_n^{1/k} = 0$  is PR on  $\mathbb{N}$  if and only if the following condition is satisfied:

# Nonlinear results/1

## Theorem (Multiplicative Rado)

A nonlinear Diophantine equation  $\prod_{i=1}^n x_i^{c_i} = 1$  is PR on  $\mathbb{N}$  if and only if the following condition is satisfied:

- “There exists a nonempty set  $J \subseteq \{1, \dots, n\}$  such that  $\sum_{j \in J} c_j = 0$ .”

Idea:  $n \rightarrow 2^n$ .

## Theorem (Lefmann)

Let  $k \in \mathbb{N}$ . A Diophantine equation of the form  $c_1 x_1^{1/k} + \dots + c_n x_n^{1/k} = 0$  is PR on  $\mathbb{N}$  if and only if the following condition is satisfied:

- “There exists a nonempty set  $J \subseteq \{1, \dots, n\}$  such that  $\sum_{j \in J} c_j = 0$ .”

# Nonlinear results/1

## Theorem (Multiplicative Rado)

A nonlinear Diophantine equation  $\prod_{i=1}^n x_i^{c_i} = 1$  is PR on  $\mathbb{N}$  if and only if the following condition is satisfied:

- “There exists a nonempty set  $J \subseteq \{1, \dots, n\}$  such that  $\sum_{j \in J} c_j = 0$ .”

Idea:  $n \rightarrow 2^n$ .

## Theorem (Lefmann)

Let  $k \in \mathbb{N}$ . A Diophantine equation of the form  $c_1 x_1^{1/k} + \dots + c_n x_n^{1/k} = 0$  is PR on  $\mathbb{N}$  if and only if the following condition is satisfied:

- “There exists a nonempty set  $J \subseteq \{1, \dots, n\}$  such that  $\sum_{j \in J} c_j = 0$ .”

Idea:  $n \rightarrow n^k$ .

## Nonlinear results/2

Theorem (Bergelson, Furstenberg, McCutcheon)

*Let  $P(z) \in \mathbb{Z}[z]$  be such that  $P(0) = 0$ . Then the equation  $x - y = P(z)$  is PR.*



## Nonlinear results/2

### Theorem (Bergelson, Furstenberg, McCutcheon)

*Let  $P(z) \in \mathbb{Z}[z]$  be such that  $P(0) = 0$ . Then the equation  $x - y = P(z)$  is PR.*

Idea: use ergodic methods (recurrence sets).

## Nonlinear results/2

### Theorem (Bergelson, Furstenberg, McCutcheon)

*Let  $P(z) \in \mathbb{Z}[z]$  be such that  $P(0) = 0$ . Then the equation  $x - y = P(z)$  is PR.*

Idea: use ergodic methods (recurrence sets).

### Theorem (Csikvári, Gyarmati and Sárközy)

*The equation  $x + y = z^2$  is not partition regular.*

## Nonlinear results/2

### Theorem (Bergelson, Furstenberg, McCutcheon)

*Let  $P(z) \in \mathbb{Z}[z]$  be such that  $P(0) = 0$ . Then the equation  $x - y = P(z)$  is PR.*

Idea: use ergodic methods (recurrence sets).

### Theorem (Csikvári, Gyarmati and Sárközy)

*The equation  $x + y = z^2$  is not partition regular.*

They asked the following question: is  $x + y = tz$  PR?

## Nonlinear results/2

### Theorem (Bergelson, Furstenberg, McCutcheon)

*Let  $P(z) \in \mathbb{Z}[z]$  be such that  $P(0) = 0$ . Then the equation  $x - y = P(z)$  is PR.*

Idea: use ergodic methods (recurrence sets).

### Theorem (Csikvári, Gyarmati and Sárközy)

*The equation  $x + y = z^2$  is not partition regular.*

They asked the following question: is  $x + y = tz$  PR?

In 2010, by using algebra in the space of ultrafilters  $\beta\mathbb{N}$ , Bergelson solved the problem in the positive.

## Nonlinear results/3

Independently, Hindman proved a more general version of Bergelson's result:

## Nonlinear results/3

Independently, Hindman proved a more general version of Bergelson's result:

### Theorem (Hindman)

*All Diophantine equations of the form  $\sum_{i=1}^n x_i = \prod_{i=1}^m y_i$  are PR.*

## Nonlinear results/3

Independently, Hindman proved a more general version of Bergelson's result:

### Theorem (Hindman)

*All Diophantine equations of the form  $\sum_{i=1}^n x_i = \prod_{i=1}^m y_i$  are PR.*

Idea: use the algebra of  $\beta\mathbb{N}$ , in particular the existence of a ultrafilter  $\mathcal{U}$  such that every set  $A \in \mathcal{U}$  is additively and multiplicatively IP.

## Nonlinear results/3

Independently, Hindman proved a more general version of Bergelson's result:

### Theorem (Hindman)

*All Diophantine equations of the form  $\sum_{i=1}^n x_i = \prod_{i=1}^m y_i$  are PR.*

Idea: use the algebra of  $\beta\mathbb{N}$ , in particular the existence of a ultrafilter  $\mathcal{U}$  such that every set  $A \in \mathcal{U}$  is additively and multiplicatively IP.

### Theorem (Luperi Baglini)

*Let  $n, m > 0$ . For every choice of sets  $F_i \subseteq \{1, \dots, m\}$ , the equation  $\sum_{i=1}^n c_i x_i (\prod_{j \in F_i} y_j) = 0$  is partition regular whenever  $\sum_{i \in J} c_j = 0$  for some nonempty  $J \subseteq \{1, \dots, m\}$ . (It is agreed that  $\prod_{j \in \emptyset} y_j = 1$ .)*



## Nonlinear results/3

Independently, Hindman proved a more general version of Bergelson's result:

### Theorem (Hindman)

*All Diophantine equations of the form  $\sum_{i=1}^n x_i = \prod_{i=1}^m y_i$  are PR.*

Idea: use the algebra of  $\beta\mathbb{N}$ , in particular the existence of a ultrafilter  $\mathcal{U}$  such that every set  $A \in \mathcal{U}$  is additively and multiplicatively IP.

### Theorem (Luperi Baglini)

*Let  $n, m > 0$ . For every choice of sets  $F_i \subseteq \{1, \dots, m\}$ , the equation  $\sum_{i=1}^n c_i x_i (\prod_{j \in F_i} y_j) = 0$  is partition regular whenever  $\sum_{i \in J} c_j = 0$  for some nonempty  $J \subseteq \{1, \dots, m\}$ . (It is agreed that  $\prod_{j \in \emptyset} y_j = 1$ .)*

Idea: use the existence of a multiplicatively idempotent ultrafilter  $\mathcal{U}$  with good linear properties; study the ultrafilter using nonstandard analysis.

## Nonlinear results/4

### Theorem (Di Nasso, Riggio)

*Let  $k, n, m \in \mathbb{N}$  be such that  $k \notin \{n, m\}$ . Then the equation  $x^m + y^n = z^k$  is not PR.*

## Nonlinear results/4

### Theorem (Di Nasso, Riggio)

*Let  $k, n, m \in \mathbb{N}$  be such that  $k \notin \{n, m\}$ . Then the equation  $x^m + y^n = z^k$  is not PR.*

Idea: use nonstandard analysis, write numbers in base  $p$  for a sufficiently large prime number  $p$ .

## Nonlinear results/4

### Theorem (Di Nasso, Riggio)

*Let  $k, n, m \in \mathbb{N}$  be such that  $k \notin \{n, m\}$ . Then the equation  $x^m + y^n = z^k$  is not PR.*

Idea: use nonstandard analysis, write numbers in base  $p$  for a sufficiently large prime number  $p$ .

### Theorem (Moreira)

*Let  $\sum_{i=1}^n c_i = 0$ . Then  $\sum_{i=1}^n c_i x_i^2 = y$  is PR.*

## Nonlinear results/4

### Theorem (Di Nasso, Riggio)

*Let  $k, n, m \in \mathbb{N}$  be such that  $k \notin \{n, m\}$ . Then the equation  $x^m + y^n = z^k$  is not PR.*

Idea: use nonstandard analysis, write numbers in base  $p$  for a sufficiently large prime number  $p$ .

### Theorem (Moreira)

*Let  $\sum_{i=1}^n c_i = 0$ . Then  $\sum_{i=1}^n c_i x_i^2 = y$  is PR.*

Idea: use ergodic methods involving the set of affinities  $\{x \rightarrow ax + b\}$ ; alternatively, use an embeddability property of piecewise syndetic sets w.r.t. arithmetic progressions.

# The algebra of $\beta\mathbb{N}$

## Definition

$\mathcal{U} \subseteq \wp(\mathbb{N})$  is an ultrafilter if:

# The algebra of $\beta\mathbb{N}$

## Definition

$\mathcal{U} \subseteq \wp(\mathbb{N})$  is an ultrafilter if:

- $\emptyset \notin \mathcal{U}$ ;

# The algebra of $\beta\mathbb{N}$

## Definition

$\mathcal{U} \subseteq \wp(\mathbb{N})$  is an ultrafilter if:

- $\emptyset \notin \mathcal{U}$ ;
- $A \in \mathcal{U}, A \subseteq B \Rightarrow B \in \mathcal{U}$ ;



# The algebra of $\beta\mathbb{N}$

## Definition

$\mathcal{U} \subseteq \wp(\mathbb{N})$  is an ultrafilter if:

- $\emptyset \notin \mathcal{U}$ ;
- $A \in \mathcal{U}, A \subseteq B \Rightarrow B \in \mathcal{U}$ ;
- $A, B \in \mathcal{U} \Rightarrow A \cap B \in \mathcal{U}$ ;

# The algebra of $\beta\mathbb{N}$

## Definition

$\mathcal{U} \subseteq \wp(\mathbb{N})$  is an ultrafilter if:

- $\emptyset \notin \mathcal{U}$ ;
- $A \in \mathcal{U}, A \subseteq B \Rightarrow B \in \mathcal{U}$ ;
- $A, B \in \mathcal{U} \Rightarrow A \cap B \in \mathcal{U}$ ;
- $\forall A \in \wp(\mathbb{N}) \quad A \in \mathcal{U} \Leftrightarrow A^c \notin \mathcal{U}$ .

# The algebra of $\beta\mathbb{N}$

## Definition

$\mathcal{U} \subseteq \wp(\mathbb{N})$  is an ultrafilter if:

- $\emptyset \notin \mathcal{U}$ ;
- $A \in \mathcal{U}, A \subseteq B \Rightarrow B \in \mathcal{U}$ ;
- $A, B \in \mathcal{U} \Rightarrow A \cap B \in \mathcal{U}$ ;
- $\forall A \in \wp(\mathbb{N}) \quad A \in \mathcal{U} \Leftrightarrow A^c \notin \mathcal{U}$ .

We let  $\beta\mathbb{N}$  denote the set of ultrafilters on  $\mathbb{N}$ .

# The algebra of $\beta\mathbb{N}$

## Definition

$\mathcal{U} \subseteq \wp(\mathbb{N})$  is an ultrafilter if:

- $\emptyset \notin \mathcal{U}$ ;
- $A \in \mathcal{U}, A \subseteq B \Rightarrow B \in \mathcal{U}$ ;
- $A, B \in \mathcal{U} \Rightarrow A \cap B \in \mathcal{U}$ ;
- $\forall A \in \wp(\mathbb{N}) \quad A \in \mathcal{U} \Leftrightarrow A^c \notin \mathcal{U}$ .

We let  $\beta\mathbb{N}$  denote the set of ultrafilters on  $\mathbb{N}$ .

Topology: for every  $A \in \wp(\mathbb{N})$  let  $\Theta_A = \{\mathcal{U} \in \beta\mathbb{N} \mid A \in \mathcal{U}\}$ .

# The algebra of $\beta\mathbb{N}$

## Definition

$\mathcal{U} \subseteq \wp(\mathbb{N})$  is an ultrafilter if:

- $\emptyset \notin \mathcal{U}$ ;
- $A \in \mathcal{U}, A \subseteq B \Rightarrow B \in \mathcal{U}$ ;
- $A, B \in \mathcal{U} \Rightarrow A \cap B \in \mathcal{U}$ ;
- $\forall A \in \wp(\mathbb{N}) \quad A \in \mathcal{U} \Leftrightarrow A^c \notin \mathcal{U}$ .

We let  $\beta\mathbb{N}$  denote the set of ultrafilters on  $\mathbb{N}$ .

Topology: for every  $A \in \wp(\mathbb{N})$  let  $\Theta_A = \{\mathcal{U} \in \beta\mathbb{N} \mid A \in \mathcal{U}\}$ .

If  $\mathcal{U}, \mathcal{V} \in \beta\mathbb{N}$  and  $A \subseteq \mathbb{N}$ , we let

$$A \in \mathcal{U} \oplus \mathcal{V} \Leftrightarrow \{n \mid A - n \in \mathcal{V}\} \in \mathcal{U},$$

# The algebra of $\beta\mathbb{N}$

## Definition

$\mathcal{U} \subseteq \wp(\mathbb{N})$  is an ultrafilter if:

- $\emptyset \notin \mathcal{U}$ ;
- $A \in \mathcal{U}, A \subseteq B \Rightarrow B \in \mathcal{U}$ ;
- $A, B \in \mathcal{U} \Rightarrow A \cap B \in \mathcal{U}$ ;
- $\forall A \in \wp(\mathbb{N}) \quad A \in \mathcal{U} \Leftrightarrow A^c \notin \mathcal{U}$ .

We let  $\beta\mathbb{N}$  denote the set of ultrafilters on  $\mathbb{N}$ .

Topology: for every  $A \in \wp(\mathbb{N})$  let  $\Theta_A = \{\mathcal{U} \in \beta\mathbb{N} \mid A \in \mathcal{U}\}$ .

If  $\mathcal{U}, \mathcal{V} \in \beta\mathbb{N}$  and  $A \subseteq \mathbb{N}$ , we let

$$A \in \mathcal{U} \oplus \mathcal{V} \Leftrightarrow \{n \mid A - n \in \mathcal{V}\} \in \mathcal{U},$$

where  $A - n = \{m \in \mathbb{N} \mid m + n \in A\}$ .

# The algebra of $\beta\mathbb{N}$

## Definition

$\mathcal{U} \subseteq \wp(\mathbb{N})$  is an ultrafilter if:

- $\emptyset \notin \mathcal{U}$ ;
- $A \in \mathcal{U}, A \subseteq B \Rightarrow B \in \mathcal{U}$ ;
- $A, B \in \mathcal{U} \Rightarrow A \cap B \in \mathcal{U}$ ;
- $\forall A \in \wp(\mathbb{N}) \quad A \in \mathcal{U} \Leftrightarrow A^c \notin \mathcal{U}$ .

We let  $\beta\mathbb{N}$  denote the set of ultrafilters on  $\mathbb{N}$ .

Topology: for every  $A \in \wp(\mathbb{N})$  let  $\Theta_A = \{\mathcal{U} \in \beta\mathbb{N} \mid A \in \mathcal{U}\}$ .

If  $\mathcal{U}, \mathcal{V} \in \beta\mathbb{N}$  and  $A \subseteq \mathbb{N}$ , we let

$$A \in \mathcal{U} \oplus \mathcal{V} \Leftrightarrow \{n \mid A - n \in \mathcal{V}\} \in \mathcal{U},$$

where  $A - n = \{m \in \mathbb{N} \mid m + n \in A\}$ . Similarly one can define  $\mathcal{U} \odot \mathcal{V}$ .

# Partition regularity as a ultrafilters problem

$\beta\mathbb{N}$  turns out to be a natural setting where to study PR problems because of the following characterization (which is here given for equations, but holds in a way more general fashion):



# Partition regularity as a ultrafilters problem

$\beta\mathbb{N}$  turns out to be a natural setting where to study PR problems because of the following characterization (which is here given for equations, but holds in a way more general fashion):

## Proposition

*A Diophantine equation  $P(x_1, \dots, x_n) = 0$  is PR if and only if there exists  $\mathcal{U} \in \beta\mathbb{N}$  such that for every  $A \in \mathcal{U}$  there exists  $a_1, \dots, a_n \in A$  with  $P(a_1, \dots, a_n) = 0$ .*

# Partition regularity as a ultrafilters problem

$\beta\mathbb{N}$  turns out to be a natural setting where to study PR problems because of the following characterization (which is here given for equations, but holds in a way more general fashion):

## Proposition

*A Diophantine equation  $P(x_1, \dots, x_n) = 0$  is PR if and only if there exists  $\mathcal{U} \in \beta\mathbb{N}$  such that for every  $A \in \mathcal{U}$  there exists  $a_1, \dots, a_n \in A$  with  $P(a_1, \dots, a_n) = 0$ .*

In this case, we say that  $\mathcal{U}$  witnesses the PR of the equation (notation:  $\mathcal{U} \models P(a_1, \dots, a_n) = 0$ ).

# Banach density and IP-sets

## Definition

Let  $A \subseteq \mathbb{N}$ . The *upper Banach density* of  $A$  is

$$BD(A) = \lim_{n \rightarrow +\infty} \sup_{m \in \mathbb{N}} \frac{|A \cap [m, m+n]|}{n+1}.$$

# Banach density and IP-sets

## Definition

Let  $A \subseteq \mathbb{N}$ . The *upper Banach density* of  $A$  is

$$BD(A) = \lim_{n \rightarrow +\infty} \sup_{m \in \mathbb{N}} \frac{|A \cap [m, m+n]|}{n+1}.$$

## Definition

Let  $G = (g_i)_{i \in \mathbb{N}}$  be an increasing sequence of natural numbers. The IP-set generated by  $G$  is the set of finite sums

$$FS(G) = FS(g_i)_{i \in \mathbb{N}} = \left\{ \sum_{j=1}^k g_{i_j} \mid i_1 < i_2 < \dots < i_k \right\}.$$

A set  $A \subseteq \mathbb{N}$  is called IP-large if it contains an IP-set. Multiplicative IP-sets and multiplicative IP-large sets are defined similarly.

# Special ultrafilters

Various kind of ultrafilters are important in this field.

---

<sup>1</sup>Well, as far as "simple" goes for the kind of ultrafilters used here.

## Special ultrafilters

Various kind of ultrafilters are important in this field. However, to keep things simple<sup>1</sup>, it suffices to know that

$$\overline{K(\odot)} \cap \overline{\mathbb{I}(\oplus)} \cap \mathcal{BD} \neq \emptyset.$$

Indeed, it contains all combinatorially rich ultrafilters.

---

<sup>1</sup>Well, as far as "simple" goes for the kind of ultrafilters used here.

## Special ultrafilters

Various kind of ultrafilters are important in this field. However, to keep things simple<sup>1</sup>, it suffices to know that

$$\overline{K(\odot)} \cap \overline{\mathbb{I}(\oplus)} \cap \mathcal{BD} \neq \emptyset.$$

Indeed, it contains all combinatorially rich ultrafilters.

Idea to keep in mind for what follows: there exist some super nice ultrafilters, whose existence will be used in the following.

---

<sup>1</sup>Well, as far as "simple" goes for the kind of ultrafilters used here.

# A surprisingly simple key Lemma

## Lemma

*Let  $\mathcal{U}$  be a common witness of the equations  $P_1(x_1, \dots, x_n) = 0$  and  $P_2(y_1, \dots, y_m) = 0$ .*



# A surprisingly simple key Lemma

## Lemma

*Let  $\mathcal{U}$  be a common witness of the equations  $P_1(x_1, \dots, x_n) = 0$  and  $P_2(y_1, \dots, y_m) = 0$ . Then  $\mathcal{U}$  is also a PR-witness of the system:*

$$\begin{cases} P_1(x_1, \dots, x_n) = 0; \\ P_2(y_1, \dots, y_m) = 0; \\ x_1 = y_1. \end{cases}$$

# A surprisingly simple key Lemma

## Lemma

Let  $\mathcal{U}$  be a common witness of the equations  $P_1(x_1, \dots, x_n) = 0$  and  $P_2(y_1, \dots, y_m) = 0$ . Then  $\mathcal{U}$  is also a PR-witness of the system:

$$\begin{cases} P_1(x_1, \dots, x_n) = 0; \\ P_2(y_1, \dots, y_m) = 0; \\ x_1 = y_1. \end{cases}$$

## Proof.

Let  $A \in \mathcal{U}$  be fixed. Let

$$\Lambda_1 = \{a \in A \mid \exists a_2, \dots, a_n \in A \text{ s.t. } P_1(a, a_2, \dots, a_n) = 0\},$$

$$\Lambda_2 = \{b \in A \mid \exists b_2, \dots, b_m \in A \text{ s.t. } P_2(b, b_2, \dots, b_m) = 0\}.$$

# A surprisingly simple key Lemma

## Lemma

Let  $\mathcal{U}$  be a common witness of the equations  $P_1(x_1, \dots, x_n) = 0$  and  $P_2(y_1, \dots, y_m) = 0$ . Then  $\mathcal{U}$  is also a PR-witness of the system:

$$\begin{cases} P_1(x_1, \dots, x_n) = 0; \\ P_2(y_1, \dots, y_m) = 0; \\ x_1 = y_1. \end{cases}$$

## Proof.

Let  $A \in \mathcal{U}$  be fixed. Let

$$\Lambda_1 = \{a \in A \mid \exists a_2, \dots, a_n \in A \text{ s.t. } P_1(a, a_2, \dots, a_n) = 0\},$$

$$\Lambda_2 = \{b \in A \mid \exists b_2, \dots, b_m \in A \text{ s.t. } P_2(b, b_2, \dots, b_m) = 0\}.$$

Notice that  $\Lambda_1, \Lambda_2 \in \mathcal{U}$ , as otherwise  $\neg(\mathcal{U} \models P_i = 0)$ .

# A surprisingly simple key Lemma

## Lemma

Let  $\mathcal{U}$  be a common witness of the equations  $P_1(x_1, \dots, x_n) = 0$  and  $P_2(y_1, \dots, y_m) = 0$ . Then  $\mathcal{U}$  is also a PR-witness of the system:

$$\begin{cases} P_1(x_1, \dots, x_n) = 0; \\ P_2(y_1, \dots, y_m) = 0; \\ x_1 = y_1. \end{cases}$$

## Proof.

Let  $A \in \mathcal{U}$  be fixed. Let

$$\Lambda_1 = \{a \in A \mid \exists a_2, \dots, a_n \in A \text{ s.t. } P_1(a, a_2, \dots, a_n) = 0\},$$

$$\Lambda_2 = \{b \in A \mid \exists b_2, \dots, b_m \in A \text{ s.t. } P_2(b, b_2, \dots, b_m) = 0\}.$$

Notice that  $\Lambda_1, \Lambda_2 \in \mathcal{U}$ , as otherwise  $\neg(\mathcal{U} \models P_i = 0)$ . Take  $\Lambda_1 \cap \Lambda_2$ .  $\square$

## Some examples

### Example

Take  $\mathcal{U} \models u - v = t^2$ .

# Some examples

## Example

Take  $\mathcal{U} \models u - v = t^2$ .

Then  $\mathcal{U}$  witnesses also of the PR of the system

$$\begin{cases} u_1 - y = x^2; \\ u_2 - z = t^2; \\ y = t. \end{cases}$$

# Some examples

## Example

Take  $\mathcal{U} \models u - v = t^2$ .

Then  $\mathcal{U}$  witnesses also of the PR of the system

$$\begin{cases} u_1 - y = x^2; \\ u_2 - z = t^2; \\ y = t. \end{cases}$$

It is readily seen that this is equivalent to the PR of the configuration  $\{x, y, z, y + x^2, z + y^2\}$  (which had already been proven by ergodic methods).

# Homogeneous equations

## Theorem

*Let  $P(x_1, \dots, x_n)$  be a homogeneous PR polynomial.*



# Homogeneous equations

## Theorem

*Let  $P(x_1, \dots, x_n)$  be a homogeneous PR polynomial. Then the set of its PR-witnesses*

$$\mathfrak{W}_P = \{\mathcal{U} \in \beta\mathbb{N} \mid \mathcal{U} \models P(x_1, \dots, x_n) = 0\}$$

*is a closed multiplicative two sided ideal.*

# Homogeneous equations

## Theorem

*Let  $P(x_1, \dots, x_n)$  be a homogeneous PR polynomial. Then the set of its PR-witnesses*

$$\mathfrak{W}_P = \{\mathcal{U} \in \beta\mathbb{N} \mid \mathcal{U} \models P(x_1, \dots, x_n) = 0\}$$

*is a closed multiplicative two sided ideal.*

## Corollary

*Let  $P(x_1, \dots, x_n)$  be a homogeneous PR polynomial. Then  $\mathcal{U} \models P(x_1, \dots, x_n) = 0$  for every  $\mathcal{U} \in \overline{K(\beta\mathbb{N}, \odot)}$ .*

# The first generalization result

## Theorem

*Let  $c(x_1 - x_2) = P(y_1, \dots, y_k)$  be a Diophantine equation where the polynomial  $P$  has no constant term and  $c \neq 0$ . If the set  $A \subseteq \mathbb{N}$  is IP-large and has positive Banach density then there exist  $\xi_1, \xi_2 \in A$  and mutually distinct  $\eta_1, \dots, \eta_k \in A$  such that  $c(\xi_1 - \xi_2) = P(\eta_1, \dots, \eta_k)$ . Moreover, if  $k = 1$  then one can take  $\xi_1 \neq \xi_2$ .*

# The first generalization result

## Theorem

Let  $c(x_1 - x_2) = P(y_1, \dots, y_k)$  be a Diophantine equation where the polynomial  $P$  has no constant term and  $c \neq 0$ . If the set  $A \subseteq \mathbb{N}$  is IP-large and has positive Banach density then there exist  $\xi_1, \xi_2 \in A$  and mutually distinct  $\eta_1, \dots, \eta_k \in A$  such that  $c(\xi_1 - \xi_2) = P(\eta_1, \dots, \eta_k)$ . Moreover, if  $k = 1$  then one can take  $\xi_1 \neq \xi_2$ .

## Definition

A polynomial with integer coefficients is called a *Rado polynomial* if it can be written in the form

$$c_1x_1 + \dots + c_nx_n + P(y_1, \dots, y_k)$$

where  $n \geq 2$ ,  $P$  has no constant term, and there exists a nonempty subset  $J \subseteq \{1, \dots, n\}$  such that  $\sum_{j \in J} c_j = 0$ .

# Generalized Rado

## Theorem

*Let*

$$R(x_1, \dots, x_n, y_1, \dots, y_k) = c_1 x_1 + \dots + c_n x_n + P(y_1, \dots, y_k)$$

*be a Rado polynomial. Then every ultrafilter  $\mathcal{U} \in \overline{K(\odot)} \cap \overline{\mathbb{I}(\oplus)} \cap \mathcal{BD}$  is a PR-witness of  $R = 0$ .*

# Generalized Rado

## Theorem

Let

$$R(x_1, \dots, x_n, y_1, \dots, y_k) = c_1x_1 + \dots + c_nx_n + P(y_1, \dots, y_k)$$

be a Rado polynomial. Then every ultrafilter  $\mathcal{U} \in \overline{K(\odot)} \cap \overline{\mathbb{I}(\oplus)} \cap \mathcal{BD}$  is a PR-witness of  $R = 0$ .

## Proof.

Consider the following system:

$$\begin{cases} c_1z + c_2x_2 + \dots + c_nx_n = 0; \\ c_1(w - x_1) = P(y_1, \dots, y_k); \\ z = w. \end{cases}$$



# Main positive result/1

## Theorem

*Let  $\mathfrak{F}$  be the family of polynomials whose PR on  $\mathbb{N}$  is witnessed by at least an ultrafilter  $\mathcal{U} \in \mathbb{I}(\odot) \cap \overline{K(\odot)} \cap \overline{\mathbb{I}(\oplus)} \cap \mathcal{BD}$ . Then  $\mathfrak{F}$  includes:*

# Main positive result/1

## Theorem

*Let  $\mathfrak{F}$  be the family of polynomials whose PR on  $\mathbb{N}$  is witnessed by at least an ultrafilter  $\mathcal{U} \in \mathbb{I}(\odot) \cap \overline{K(\odot)} \cap \overline{\mathbb{I}(\oplus)} \cap \mathcal{BD}$ . Then  $\mathfrak{F}$  includes:*

- *Every Rado polynomial;*



# Main positive result/1

## Theorem

Let  $\mathfrak{F}$  be the family of polynomials whose PR on  $\mathbb{N}$  is witnessed by at least an ultrafilter  $\mathcal{U} \in \mathbb{I}(\odot) \cap \overline{K(\odot)} \cap \overline{\mathbb{I}(\oplus)} \cap \mathcal{BD}$ . Then  $\mathfrak{F}$  includes:

- Every Rado polynomial;
- Every polynomial of the form

$$\sum_{i=1}^n c_i x_i \left( \prod_{j \in F_i} y_j \right)$$

where  $\sum_{i=1}^n c_i x_i$  is a Rado polynomial and sets  $F_i \subseteq \{1, \dots, m\}$ ;

# Main positive result/1

## Theorem

Let  $\mathfrak{F}$  be the family of polynomials whose PR on  $\mathbb{N}$  is witnessed by at least an ultrafilter  $\mathcal{U} \in \mathbb{I}(\odot) \cap \overline{K(\odot)} \cap \overline{\mathbb{I}(\oplus)} \cap \mathcal{BD}$ . Then  $\mathfrak{F}$  includes:

- Every Rado polynomial;
- Every polynomial of the form

$$\sum_{i=1}^n c_i x_i \left( \prod_{j \in F_i} y_j \right)$$

where  $\sum_{i=1}^n c_i x_i$  is a Rado polynomial and sets  $F_i \subseteq \{1, \dots, m\}$ ;

- Every polynomial

$$P(x, y_1, \dots, y_k) = x - \prod_{i=1}^k y_i;$$

# Main positive result/1

## Theorem

Let  $\mathfrak{F}$  be the family of polynomials whose PR on  $\mathbb{N}$  is witnessed by at least an ultrafilter  $\mathcal{U} \in \mathbb{I}(\odot) \cap \overline{K(\odot)} \cap \overline{\mathbb{I}(\oplus)} \cap \mathcal{BD}$ . Then  $\mathfrak{F}$  includes:

- Every Rado polynomial;
- Every polynomial of the form

$$\sum_{i=1}^n c_i x_i \left( \prod_{j \in F_i} y_j \right)$$

where  $\sum_{i=1}^n c_i x_i$  is a Rado polynomial and sets  $F_i \subseteq \{1, \dots, m\}$ ;

- Every polynomial

$$P(x, y_1, \dots, y_k) = x - \prod_{i=1}^k y_i;$$

# Main positive result/2

## Theorem

- *Every polynomial*

$$P(x, y_1, \dots, y_k) = x - \prod_{i=1}^k y_i^{a_i},$$

*whenever the exponents  $a_i \in \mathbb{Z}$  satisfy  $\sum_{i=1}^n a_i = 1$ .*

# Main positive result/2

## Theorem

- *Every polynomial*

$$P(x, y_1, \dots, y_k) = x - \prod_{i=1}^k y_i^{a_i},$$

*whenever the exponents  $a_i \in \mathbb{Z}$  satisfy  $\sum_{i=1}^n a_i = 1$ .*

*Moreover, the family  $\mathfrak{F}$  satisfies the following closure properties:*

# Main positive result/2

## Theorem

- *Every polynomial*

$$P(x, y_1, \dots, y_k) = x - \prod_{i=1}^k y_i^{a_i},$$

*whenever the exponents  $a_i \in \mathbb{Z}$  satisfy  $\sum_{i=1}^n a_i = 1$ .*

*Moreover, the family  $\mathfrak{F}$  satisfies the following closure properties:*

- (i) *If  $P(z, y_1, \dots, y_k) \in \mathfrak{F}$  and  $z - g(x_1, \dots, x_n) \in \mathfrak{F}$ , then  $P(g(x_1, \dots, x_n), y_1, \dots, y_k) \in \mathfrak{F}$ ;*

## Main positive result/2

### Theorem

- *Every polynomial*

$$P(x, y_1, \dots, y_k) = x - \prod_{i=1}^k y_i^{a_i},$$

whenever the exponents  $a_i \in \mathbb{Z}$  satisfy  $\sum_{i=1}^k a_i = 1$ .

Moreover, the family  $\mathfrak{F}$  satisfies the following closure properties:

- (i) If  $P(z, y_1, \dots, y_k) \in \mathfrak{F}$  and  $z - g(x_1, \dots, x_n) \in \mathfrak{F}$ , then  $P(g(x_1, \dots, x_n), y_1, \dots, y_k) \in \mathfrak{F}$ ;
- (ii) if  $P(x_1, \dots, x_n) \in \mathfrak{F}$  is homogeneous, then  $P\left(\frac{1}{x_1}, \dots, \frac{1}{x_n}\right) \in \mathfrak{F}$ .

# Some examples



## Some examples

### Example

Let  $n, m \in \mathbb{N}$  and assume that, for every  $i \leq n, j \leq m$ , the equations

$$x_{i,1} = \sum_{h=1}^{r_i} c_{i,h} x_{i,h}, \quad y_{j,1} = \sum_{k=1}^{s_j} d_{j,k} y_{j,k}$$

are PR.

## Some examples

### Example

Let  $n, m \in \mathbb{N}$  and assume that, for every  $i \leq n, j \leq m$ , the equations

$$x_{i,1} = \sum_{h=1}^{r_i} c_{i,h} x_{i,h}, \quad y_{j,1} = \sum_{k=1}^{s_j} d_{j,k} y_{j,k}$$

are PR.

Let  $a_1, \dots, a_n, b_1, \dots, b_m$  be such that  $\sum_{i=1}^n a_i = \sum_{j=1}^m b_j$  and consider the homogeneous PR equation  $\prod_{i=1}^n t_i^{a_i} = \prod_{j=1}^m z_j^{b_j}$ .

## Some examples

### Example

Let  $n, m \in \mathbb{N}$  and assume that, for every  $i \leq n, j \leq m$ , the equations

$$x_{i,1} = \sum_{h=1}^{r_i} c_{i,h} x_{i,h}, \quad y_{j,1} = \sum_{k=1}^{s_j} d_{j,k} y_{j,k}$$

are PR.

Let  $a_1, \dots, a_n, b_1, \dots, b_m$  be such that  $\sum_{i=1}^n a_i = \sum_{j=1}^m b_j$  and consider the homogeneous PR equation  $\prod_{i=1}^n t_i^{a_i} = \prod_{j=1}^m z_j^{b_j}$ .

All these equations are PR and homogeneous and therefore, by the closure property (i), also

$$\prod_{i=1}^n \left( \sum_{h=1}^{r_i} c_{i,h} x_{i,h} \right)^{a_i} = \prod_{j=1}^m \left( \sum_{k=1}^{s_j} d_{j,k} y_{j,k} \right)^{b_j}$$

is PR.

## Some examples

### Example

Let  $n, m \in \mathbb{N}$  and assume that, for every  $i \leq n$ ,  $j \leq m$ , the equations

$$x_{i,1} = \sum_{h=1}^{r_i} c_{i,h} x_{i,h}, \quad y_{j,1} = \sum_{k=1}^{s_j} d_{j,k} y_{j,k}$$

are PR.

Let  $a_1, \dots, a_n, b_1, \dots, b_m$  be such that  $\sum_{i=1}^n a_i = \sum_{j=1}^m b_j$  and consider the homogeneous PR equation  $\prod_{i=1}^n t_i^{a_i} = \prod_{j=1}^m z_j^{b_j}$ .

All these equations are PR and homogeneous and therefore, by the closure property (i), also

$$\prod_{i=1}^n \left( \sum_{h=1}^{r_i} c_{i,h} x_{i,h} \right)^{a_i} = \prod_{j=1}^m \left( \sum_{k=1}^{s_j} d_{j,k} y_{j,k} \right)^{b_j}$$

is PR.

## Some examples

### Example

For every  $n \in \mathbb{N}$ , the polynomial  $u - v - z^n$  is in  $\mathfrak{F}$ ;

## Some examples

### Example

For every  $n \in \mathbb{N}$ , the polynomial  $u - v - z^n$  is in  $\mathfrak{F}$ ; moreover, for every  $k \geq 2$  the function  $x = \prod_{j=1}^k x_j$  is in  $\mathfrak{F}$ .

## Some examples

### Example

For every  $n \in \mathbb{N}$ , the polynomial  $u - v - z^n$  is in  $\mathfrak{F}$ ; moreover, for every  $k \geq 2$  the function  $x = \prod_{j=1}^k x_j$  is in  $\mathfrak{F}$ . Therefore, for every  $h, k \geq 2$  we can apply the closure property (i) of  $\mathfrak{F}$  to the system

$$\begin{cases} u - v = z^n; \\ x = \prod_{j=1}^h x_j; \\ y = \prod_{j=1}^k y_j; \\ x = t, y = v. \end{cases}$$

## Some examples

### Example

For every  $n \in \mathbb{N}$ , the polynomial  $u - v - z^n$  is in  $\mathfrak{F}$ ; moreover, for every  $k \geq 2$  the function  $x = \prod_{j=1}^k x_j$  is in  $\mathfrak{F}$ . Therefore, for every  $h, k \geq 2$  we can apply the closure property (i) of  $\mathfrak{F}$  to the system

$$\begin{cases} u - v = z^n; \\ x = \prod_{j=1}^h x_j; \\ y = \prod_{j=1}^k y_j; \\ x = t, y = v. \end{cases}$$

Hence  $\prod_{j=1}^h x_j - \prod_{j=1}^k y_j = z^n$  is in  $\mathfrak{F}$ . In particular,  $x_1 x_2 - y_1 y_2 = z^2$  is PR.



## Some examples

### Example

For every  $n \in \mathbb{N}$ , the polynomial  $u - v - z^n$  is in  $\mathfrak{F}$ ; moreover, for every  $k \geq 2$  the function  $x = \prod_{j=1}^k x_j$  is in  $\mathfrak{F}$ . Therefore, for every  $h, k \geq 2$  we can apply the closure property (i) of  $\mathfrak{F}$  to the system

$$\begin{cases} u - v = z^n; \\ x = \prod_{j=1}^h x_j; \\ y = \prod_{j=1}^k y_j; \\ x = t, y = v. \end{cases}$$

Hence  $\prod_{j=1}^h x_j - \prod_{j=1}^k y_j = z^n$  is in  $\mathfrak{F}$ . In particular,  $x_1 x_2 - y_1 y_2 = z^2$  is PR.

### Example

$P(x_1, x_2, x_3) = x_1 x_2 - 2x_3$  is PR but it does not belong to  $\mathfrak{F}$ .

# Nonstandard analysis: basic idea

Nonstandard analysis essentially consists of two properties:

# Nonstandard analysis: basic idea

Nonstandard analysis essentially consists of two properties:

- 1 Every mathematical object  $X$  is extended to an object  ${}^*X$ , called its hyper-extension or nonstandard extension.

# Nonstandard analysis: basic idea

Nonstandard analysis essentially consists of two properties:

- ① Every mathematical object  $X$  is extended to an object  ${}^*X$ , called its hyper-extension or nonstandard extension.
- ②  ${}^*X$  is a sort of weakly isomorphic copy of  $X$ , in the sense that it satisfies the same **elementary properties** as  $X$ .

# Nonstandard analysis: basic idea

Nonstandard analysis essentially consists of two properties:

- 1 Every mathematical object  $X$  is extended to an object  ${}^*X$ , called its hyper-extension or nonstandard extension.
- 2  ${}^*X$  is a sort of weakly isomorphic copy of  $X$ , in the sense that it satisfies the same **elementary properties** as  $X$ .

A property is elementary if it talks about elements of  $X$  (it is not elementary when talks about subsets or functions).

# Nonstandard analysis: basic idea

Nonstandard analysis essentially consists of two properties:

- 1 Every mathematical object  $X$  is extended to an object  ${}^*X$ , called its hyper-extension or nonstandard extension.
- 2  ${}^*X$  is a sort of weakly isomorphic copy of  $X$ , in the sense that it satisfies the same **elementary properties** as  $X$ .

A property is elementary if it talks about elements of  $X$  (it is not elementary when talks about subsets or functions).

The preservation of elementary properties when taking hyper-extensions is called **transfer principle**.

## $u$ -equivalence and partition regularity

\* $\mathbb{N}$  is the hyper-extension of  $\mathbb{N}$ .

## *u*-equivalence and partition regularity

${}^*\mathbb{N}$  is the hyper-extension of  $\mathbb{N}$ . Its elements have the same elementary properties of natural numbers:



## $u$ -equivalence and partition regularity

${}^*\mathbb{N}$  is the hyper-extension of  $\mathbb{N}$ . Its elements have the same elementary properties of natural numbers:

- addition can be extended so to make  ${}^*\mathbb{N}$  a semigroup;

## $u$ -equivalence and partition regularity

${}^*\mathbb{N}$  is the hyper-extension of  $\mathbb{N}$ . Its elements have the same elementary properties of natural numbers:

- addition can be extended so to make  ${}^*\mathbb{N}$  a semigroup;
- the same with multiplication;

## $u$ -equivalence and partition regularity

${}^*\mathbb{N}$  is the hyper-extension of  $\mathbb{N}$ . Its elements have the same elementary properties of natural numbers:

- addition can be extended so to make  ${}^*\mathbb{N}$  a semigroup;
- the same with multiplication;
- the linear order of  $\mathbb{N}$  can be extended to a linear order of  ${}^*\mathbb{N}$ ;

## $u$ -equivalence and partition regularity

${}^*\mathbb{N}$  is the hyper-extension of  $\mathbb{N}$ . Its elements have the same elementary properties of natural numbers:

- addition can be extended so to make  ${}^*\mathbb{N}$  a semigroup;
- the same with multiplication;
- the linear order of  $\mathbb{N}$  can be extended to a linear order of  ${}^*\mathbb{N}$ ;
- and so on.

## $u$ -equivalence and partition regularity

${}^*\mathbb{N}$  is the hyper-extension of  $\mathbb{N}$ . Its elements have the same elementary properties of natural numbers:

- addition can be extended so to make  ${}^*\mathbb{N}$  a semigroup;
- the same with multiplication;
- the linear order of  $\mathbb{N}$  can be extended to a linear order of  ${}^*\mathbb{N}$ ;
- and so on.

In  ${}^*\mathbb{N}$  we have **infinite numbers**, e.g. elements  $\alpha \in {}^*\mathbb{N}$  such that  $\alpha > n$  for every  $n \in \mathbb{N}$ .

## $u$ -equivalence and partition regularity

${}^*\mathbb{N}$  is the hyper-extension of  $\mathbb{N}$ . Its elements have the same elementary properties of natural numbers:

- addition can be extended so to make  ${}^*\mathbb{N}$  a semigroup;
- the same with multiplication;
- the linear order of  $\mathbb{N}$  can be extended to a linear order of  ${}^*\mathbb{N}$ ;
- and so on.

In  ${}^*\mathbb{N}$  we have **infinite numbers**, e.g. elements  $\alpha \in {}^*\mathbb{N}$  such that  $\alpha > n$  for every  $n \in \mathbb{N}$ .

Example of non-elementary property: the well-order. In fact, the set of infinite elements does not have a minimum.

## $u$ -equivalence and partition regularity

### Definition

Two hypernatural numbers  $\xi, \xi' \in {}^*\mathbb{N}$  are  *$u$ -equivalent* if they cannot be distinguished by any hyper-extension, i.e. if for every  $A \subseteq \mathbb{N}$  one has either  $\xi, \xi' \in {}^*A$  or  $\xi, \xi' \notin {}^*A$ .

## $u$ -equivalence and partition regularity

### Definition

*Two hypernatural numbers  $\xi, \xi' \in {}^*\mathbb{N}$  are  $u$ -equivalent if they cannot be distinguished by any hyper-extension, i.e. if for every  $A \subseteq \mathbb{N}$  one has either  $\xi, \xi' \in {}^*A$  or  $\xi, \xi' \notin {}^*A$ .*

Ultrafilters and hypernaturals can be identified:



## $u$ -equivalence and partition regularity

### Definition

Two hypernatural numbers  $\xi, \xi' \in {}^*\mathbb{N}$  are  *$u$ -equivalent* if they cannot be distinguished by any hyper-extension, i.e. if for every  $A \subseteq \mathbb{N}$  one has either  $\xi, \xi' \in {}^*A$  or  $\xi, \xi' \notin {}^*A$ .

Ultrafilters and hypernaturals can be identified:

- $\alpha \rightarrow \mathcal{U}_\alpha = \{A \in \wp(\mathbb{N}) \mid \alpha \in {}^*A\}$ ;

## $u$ -equivalence and partition regularity

### Definition

Two hypernatural numbers  $\xi, \xi' \in {}^*\mathbb{N}$  are  *$u$ -equivalent* if they cannot be distinguished by any hyper-extension, i.e. if for every  $A \subseteq \mathbb{N}$  one has either  $\xi, \xi' \in {}^*A$  or  $\xi, \xi' \notin {}^*A$ .

Ultrafilters and hypernaturals can be identified:

- $\alpha \rightarrow \mathcal{U}_\alpha = \{A \in \wp(\mathbb{N}) \mid \alpha \in {}^*A\}$ ;
- $\mathcal{U} \rightarrow \mu(\mathcal{U}) = \{\alpha \in {}^*\mathbb{N} \mid \mathcal{U} = \mathcal{U}_\alpha\}$ .

# $u$ -equivalence and partition regularity

## Definition

Two hypernatural numbers  $\xi, \xi' \in {}^*\mathbb{N}$  are  *$u$ -equivalent* if they cannot be distinguished by any hyper-extension, i.e. if for every  $A \subseteq \mathbb{N}$  one has either  $\xi, \xi' \in {}^*A$  or  $\xi, \xi' \notin {}^*A$ .

Ultrafilters and hypernaturals can be identified:

- $\alpha \rightarrow \mathcal{U}_\alpha = \{A \in \wp(\mathbb{N}) \mid \alpha \in {}^*A\}$ ;
- $\mathcal{U} \rightarrow \mu(\mathcal{U}) = \{\alpha \in {}^*\mathbb{N} \mid \mathcal{U} = \mathcal{U}_\alpha\}$ .

## Proposition

A Diophantine equation  $P(x_1, \dots, x_n) = 0$  is PR if and only if there exist  $u$ -equivalent hypernatural numbers  $\xi_1, \dots, \xi_n$  with  ${}^*P(\xi_1, \dots, \xi_n) = 0$ .

## Multi-index notations

- An  $n$ -dimensional **multi-index** is an  $n$ -tuple  $\alpha = (\alpha_1, \dots, \alpha_n) \in \mathbb{N}_0^n$ ;

## Multi-index notations

- An  $n$ -dimensional **multi-index** is an  $n$ -tuple  $\alpha = (\alpha_1, \dots, \alpha_n) \in \mathbb{N}_0^n$ ;
- $\alpha \leq \beta$  means that  $\alpha_i \leq \beta_i$  for all  $i = 1, \dots, n$ ;

## Multi-index notations

- An  $n$ -dimensional **multi-index** is an  $n$ -tuple  $\alpha = (\alpha_1, \dots, \alpha_n) \in \mathbb{N}_0^n$ ;
- $\alpha \leq \beta$  means that  $\alpha_i \leq \beta_i$  for all  $i = 1, \dots, n$ ;
- $\alpha < \beta$  means that  $\alpha \leq \beta$  and  $\alpha \neq \beta$ ;

## Multi-index notations

- An  $n$ -dimensional **multi-index** is an  $n$ -tuple  $\alpha = (\alpha_1, \dots, \alpha_n) \in \mathbb{N}_0^n$ ;
- $\alpha \leq \beta$  means that  $\alpha_i \leq \beta_i$  for all  $i = 1, \dots, n$ ;
- $\alpha < \beta$  means that  $\alpha \leq \beta$  and  $\alpha \neq \beta$ ;
- If  $\mathbf{x} = (x_1, \dots, x_n)$  is vector and  $\alpha = (\alpha_1, \dots, \alpha_n)$  is a multi-index, the product  $\prod_{i=1}^n x_i^{\alpha_i}$  is denoted by  $\mathbf{x}^\alpha$ ;

## Multi-index notations

- An  $n$ -dimensional **multi-index** is an  $n$ -tuple  $\alpha = (\alpha_1, \dots, \alpha_n) \in \mathbb{N}_0^n$ ;
- $\alpha \leq \beta$  means that  $\alpha_i \leq \beta_i$  for all  $i = 1, \dots, n$ ;
- $\alpha < \beta$  means that  $\alpha \leq \beta$  and  $\alpha \neq \beta$ ;
- If  $\mathbf{x} = (x_1, \dots, x_n)$  is vector and  $\alpha = (\alpha_1, \dots, \alpha_n)$  is a multi-index, the product  $\prod_{i=1}^n x_i^{\alpha_i}$  is denoted by  $\mathbf{x}^\alpha$ ;
- The **length** of a multi-index  $\alpha = (\alpha_1, \dots, \alpha_n)$  is  $|\alpha| = \sum_{i=1}^n \alpha_i$ ;



## Multi-index notations

- An  $n$ -dimensional **multi-index** is an  $n$ -tuple  $\alpha = (\alpha_1, \dots, \alpha_n) \in \mathbb{N}_0^n$ ;
- $\alpha \leq \beta$  means that  $\alpha_i \leq \beta_i$  for all  $i = 1, \dots, n$ ;
- $\alpha < \beta$  means that  $\alpha \leq \beta$  and  $\alpha \neq \beta$ ;
- If  $\mathbf{x} = (x_1, \dots, x_n)$  is vector and  $\alpha = (\alpha_1, \dots, \alpha_n)$  is a multi-index, the product  $\prod_{i=1}^n x_i^{\alpha_i}$  is denoted by  $\mathbf{x}^\alpha$ ;
- The **length** of a multi-index  $\alpha = (\alpha_1, \dots, \alpha_n)$  is  $|\alpha| = \sum_{i=1}^n \alpha_i$ ;
- A set  $I$  of  $n$ -dimensional multi-indexes having all the same length is called *homogeneous*;

## Multi-index notations

- An  $n$ -dimensional **multi-index** is an  $n$ -tuple  $\alpha = (\alpha_1, \dots, \alpha_n) \in \mathbb{N}_0^n$ ;
- $\alpha \leq \beta$  means that  $\alpha_i \leq \beta_i$  for all  $i = 1, \dots, n$ ;
- $\alpha < \beta$  means that  $\alpha \leq \beta$  and  $\alpha \neq \beta$ ;
- If  $\mathbf{x} = (x_1, \dots, x_n)$  is vector and  $\alpha = (\alpha_1, \dots, \alpha_n)$  is a multi-index, the product  $\prod_{i=1}^n x_i^{\alpha_i}$  is denoted by  $\mathbf{x}^\alpha$ ;
- The **length** of a multi-index  $\alpha = (\alpha_1, \dots, \alpha_n)$  is  $|\alpha| = \sum_{i=1}^n \alpha_i$ ;
- A set  $I$  of  $n$ -dimensional multi-indexes having all the same length is called *homogeneous*;
- Polynomials  $P \in \mathbb{Z}[x_1, \dots, x_n]$  are written in the form  $P(\mathbf{x}) = \sum_{\alpha} c_{\alpha} \mathbf{x}^{\alpha}$  where  $\alpha$  are multi-indexes;

## Multi-index notations

- An  $n$ -dimensional **multi-index** is an  $n$ -tuple  $\alpha = (\alpha_1, \dots, \alpha_n) \in \mathbb{N}_0^n$ ;
- $\alpha \leq \beta$  means that  $\alpha_i \leq \beta_i$  for all  $i = 1, \dots, n$ ;
- $\alpha < \beta$  means that  $\alpha \leq \beta$  and  $\alpha \neq \beta$ ;
- If  $\mathbf{x} = (x_1, \dots, x_n)$  is vector and  $\alpha = (\alpha_1, \dots, \alpha_n)$  is a multi-index, the product  $\prod_{i=1}^n x_i^{\alpha_i}$  is denoted by  $\mathbf{x}^\alpha$ ;
- The **length** of a multi-index  $\alpha = (\alpha_1, \dots, \alpha_n)$  is  $|\alpha| = \sum_{i=1}^n \alpha_i$ ;
- A set  $I$  of  $n$ -dimensional multi-indexes having all the same length is called *homogeneous*;
- Polynomials  $P \in \mathbb{Z}[x_1, \dots, x_n]$  are written in the form  $P(\mathbf{x}) = \sum_{\alpha} c_{\alpha} \mathbf{x}^{\alpha}$  where  $\alpha$  are multi-indexes;
- The **support** of  $P$  is the finite set  $\text{supp}(P) = \{\alpha \mid c_{\alpha} \neq 0\}$ ;

## Multi-index notations

- An  $n$ -dimensional **multi-index** is an  $n$ -tuple  $\alpha = (\alpha_1, \dots, \alpha_n) \in \mathbb{N}_0^n$ ;
- $\alpha \leq \beta$  means that  $\alpha_i \leq \beta_i$  for all  $i = 1, \dots, n$ ;
- $\alpha < \beta$  means that  $\alpha \leq \beta$  and  $\alpha \neq \beta$ ;
- If  $\mathbf{x} = (x_1, \dots, x_n)$  is vector and  $\alpha = (\alpha_1, \dots, \alpha_n)$  is a multi-index, the product  $\prod_{i=1}^n x_i^{\alpha_i}$  is denoted by  $\mathbf{x}^\alpha$ ;
- The **length** of a multi-index  $\alpha = (\alpha_1, \dots, \alpha_n)$  is  $|\alpha| = \sum_{i=1}^n \alpha_i$ ;
- A set  $I$  of  $n$ -dimensional multi-indexes having all the same length is called *homogeneous*;
- Polynomials  $P \in \mathbb{Z}[x_1, \dots, x_n]$  are written in the form  $P(\mathbf{x}) = \sum_{\alpha} c_{\alpha} \mathbf{x}^{\alpha}$  where  $\alpha$  are multi-indexes;
- The **support** of  $P$  is the finite set  $\text{supp}(P) = \{\alpha \mid c_{\alpha} \neq 0\}$ ;
- A polynomial  $P(\mathbf{x}) = \sum_{\alpha} c_{\alpha} \mathbf{x}^{\alpha}$  is **homogeneous** if  $\text{supp}(P)$  is a homogeneous set of indexes.

# Minimal and maximal indices

## Definition

Let  $P(\mathbf{x}) = \sum_{\alpha} c_{\alpha} \mathbf{x}^{\alpha} \in \mathbb{Z}[x_1, \dots, x_n]$ . We say that a multi-index  $\alpha \in \text{supp}(P)$  is *minimal* if there are no  $\beta \in \text{supp}(P)$  with  $\beta < \alpha$ . The notion of *maximal* multi-index is defined similarly.

# Minimal and maximal indices

## Definition

Let  $P(\mathbf{x}) = \sum_{\alpha} c_{\alpha} \mathbf{x}^{\alpha} \in \mathbb{Z}[x_1, \dots, x_n]$ . We say that a multi-index  $\alpha \in \text{supp}(P)$  is *minimal* if there are no  $\beta \in \text{supp}(P)$  with  $\beta < \alpha$ . The notion of *maximal* multi-index is defined similarly. A nonempty set  $J \subseteq \text{supp}(P)$  is called a *Rado set of indexes* if for every  $\alpha, \beta \in J$  there exists a nonempty  $\Lambda \subseteq \{1, \dots, n\}$  with  $\sum_{i \in \Lambda} \alpha_i = \sum_{i \in \Lambda} \beta_i$ .

# Minimal and maximal indices

## Definition

Let  $P(\mathbf{x}) = \sum_{\alpha} c_{\alpha} \mathbf{x}^{\alpha} \in \mathbb{Z}[x_1, \dots, x_n]$ . We say that a multi-index  $\alpha \in \text{supp}(P)$  is *minimal* if there are no  $\beta \in \text{supp}(P)$  with  $\beta < \alpha$ . The notion of *maximal* multi-index is defined similarly. A nonempty set  $J \subseteq \text{supp}(P)$  is called a *Rado set of indexes* if for every  $\alpha, \beta \in J$  there exists a nonempty  $\Lambda \subseteq \{1, \dots, n\}$  with  $\sum_{i \in \Lambda} \alpha_i = \sum_{i \in \Lambda} \beta_i$ .

For linear polynomials, every nonempty  $J \subseteq \text{Supp}(P) = \{\alpha(1), \dots, \alpha(n)\}$  is a Rado set of both minimal and maximal indices.

# Minimal and maximal indices

## Definition

Let  $P(\mathbf{x}) = \sum_{\alpha} c_{\alpha} \mathbf{x}^{\alpha} \in \mathbb{Z}[x_1, \dots, x_n]$ . We say that a multi-index  $\alpha \in \text{supp}(P)$  is *minimal* if there are no  $\beta \in \text{supp}(P)$  with  $\beta < \alpha$ . The notion of *maximal* multi-index is defined similarly. A nonempty set  $J \subseteq \text{supp}(P)$  is called a *Rado set of indexes* if for every  $\alpha, \beta \in J$  there exists a nonempty  $\Lambda \subseteq \{1, \dots, n\}$  with  $\sum_{i \in \Lambda} \alpha_i = \sum_{i \in \Lambda} \beta_i$ .

For linear polynomials, every nonempty  $J \subseteq \text{Supp}(P) = \{\alpha(1), \dots, \alpha(n)\}$  is a Rado set of both minimal and maximal indices.

## Example

In  $c_1 x_1^2 x_2 x_3 + c_2 x_1 x_2^2 x_3^7 + c_3 x_1^2 x_2^2 x_3^2 x_4$ , the set  $J = \{1, 2\}$  is a Rado set of minimal (but not maximal) indices.



## General necessary condition

### Theorem

*Let  $P(\mathbf{x}) = \sum_{\alpha} c_{\alpha} \mathbf{x}^{\alpha} \in \mathbb{Z}[x_1, \dots, x_n]$  be a polynomial with no constant term.*

## General necessary condition

### Theorem

*Let  $P(\mathbf{x}) = \sum_{\alpha} c_{\alpha} \mathbf{x}^{\alpha} \in \mathbb{Z}[x_1, \dots, x_n]$  be a polynomial with no constant term. Suppose there exists a prime  $p$  such that:*

# General necessary condition

## Theorem

Let  $P(\mathbf{x}) = \sum_{\alpha} c_{\alpha} \mathbf{x}^{\alpha} \in \mathbb{Z}[x_1, \dots, x_n]$  be a polynomial with no constant term. Suppose there exists a prime  $p$  such that:

- 1  $\sum_{\alpha} c_{\alpha} z^{|\alpha|} \equiv 0 \pmod{p}$  has no solutions  $z \not\equiv 0$ ;

# General necessary condition

## Theorem

Let  $P(\mathbf{x}) = \sum_{\alpha} c_{\alpha} \mathbf{x}^{\alpha} \in \mathbb{Z}[x_1, \dots, x_n]$  be a polynomial with no constant term. Suppose there exists a prime  $p$  such that:

- 1  $\sum_{\alpha} c_{\alpha} z^{|\alpha|} \equiv 0 \pmod{p}$  has no solutions  $z \neq 0$ ;
- 2 For every Rado set  $J$  of minimal indexes,  $\sum_{\alpha \in J} c_{\alpha} z^{|\alpha|} \equiv 0 \pmod{p}$  has no solutions  $z \neq 0$ .

# General necessary condition

## Theorem

Let  $P(\mathbf{x}) = \sum_{\alpha} c_{\alpha} \mathbf{x}^{\alpha} \in \mathbb{Z}[x_1, \dots, x_n]$  be a polynomial with no constant term. Suppose there exists a prime  $p$  such that:

- 1  $\sum_{\alpha} c_{\alpha} z^{|\alpha|} \equiv 0 \pmod{p}$  has no solutions  $z \not\equiv 0$ ;
- 2 For every Rado set  $J$  of minimal indexes,  $\sum_{\alpha \in J} c_{\alpha} z^{|\alpha|} \equiv 0 \pmod{p}$  has no solutions  $z \not\equiv 0$ .

Then  $P(\mathbf{x})$  is not PR, except possibly for constant solutions  $x_1 = \dots = x_n$ .

# General necessary condition

## Theorem

Let  $P(\mathbf{x}) = \sum_{\alpha} c_{\alpha} \mathbf{x}^{\alpha} \in \mathbb{Z}[x_1, \dots, x_n]$  be a polynomial with no constant term. Suppose there exists a prime  $p$  such that:

- 1  $\sum_{\alpha} c_{\alpha} z^{|\alpha|} \equiv 0 \pmod{p}$  has no solutions  $z \neq 0$ ;
- 2 For every Rado set  $J$  of minimal indexes,  $\sum_{\alpha \in J} c_{\alpha} z^{|\alpha|} \equiv 0 \pmod{p}$  has no solutions  $z \neq 0$ .

Then  $P(\mathbf{x})$  is not PR, except possibly for constant solutions  $x_1 = \dots = x_n$ .

## Proof.

Pick infinite  $\xi_1 \underset{\sim}{\sim} \dots \underset{\sim}{\sim} \xi_n$  such that  $P(\boldsymbol{\xi}) = \sum_{\alpha} c_{\alpha} \boldsymbol{\xi}^{\alpha} = 0$ .

# General necessary condition

## Theorem

Let  $P(\mathbf{x}) = \sum_{\alpha} c_{\alpha} \mathbf{x}^{\alpha} \in \mathbb{Z}[x_1, \dots, x_n]$  be a polynomial with no constant term. Suppose there exists a prime  $p$  such that:

- 1  $\sum_{\alpha} c_{\alpha} z^{|\alpha|} \equiv 0 \pmod{p}$  has no solutions  $z \neq 0$ ;
- 2 For every Rado set  $J$  of minimal indexes,  $\sum_{\alpha \in J} c_{\alpha} z^{|\alpha|} \equiv 0 \pmod{p}$  has no solutions  $z \neq 0$ .

Then  $P(\mathbf{x})$  is not PR, except possibly for constant solutions  $x_1 = \dots = x_n$ .

## Proof.

Pick infinite  $\xi_1 \underset{\sim}{\sim} \dots \underset{\sim}{\sim} \xi_n$  such that  $P(\boldsymbol{\xi}) = \sum_{\alpha} c_{\alpha} \boldsymbol{\xi}^{\alpha} = 0$ . Write  $\xi_i$  in base  $p$ .

# General necessary condition

## Theorem

Let  $P(\mathbf{x}) = \sum_{\alpha} c_{\alpha} \mathbf{x}^{\alpha} \in \mathbb{Z}[x_1, \dots, x_n]$  be a polynomial with no constant term. Suppose there exists a prime  $p$  such that:

- 1  $\sum_{\alpha} c_{\alpha} z^{|\alpha|} \equiv 0 \pmod{p}$  has no solutions  $z \neq 0$ ;
- 2 For every Rado set  $J$  of minimal indexes,  $\sum_{\alpha \in J} c_{\alpha} z^{|\alpha|} \equiv 0 \pmod{p}$  has no solutions  $z \neq 0$ .

Then  $P(\mathbf{x})$  is not PR, except possibly for constant solutions  $x_1 = \dots = x_n$ .

## Proof.

Pick infinite  $\xi_1 \approx \dots \approx \xi_n$  such that  $P(\boldsymbol{\xi}) = \sum_{\alpha} c_{\alpha} \boldsymbol{\xi}^{\alpha} = 0$ . Write  $\xi_i$  in base  $p$ . Find the absurd playing with the exponents and the coefficients in this expansion.  $\square$



# Examples

# Examples

## Example

Let  $P(x_1, x_2, x_3) = x_1^2 x_2 - 2x_3$ . Pick any prime number  $p$  with  $p \equiv 3$  or  $p \equiv 5 \pmod{8}$ , so that 2 is not a quadratic residue modulo  $p$ .

# Examples

## Example

Let  $P(x_1, x_2, x_3) = x_1^2 x_2 - 2x_3$ . Pick any prime number  $p$  with  $p \equiv 3$  or  $p \equiv 5 \pmod{8}$ , so that 2 is not a quadratic residue modulo  $p$ . Then condition (1) is satisfied because  $z^3 - 2z \equiv 0$  iff  $z \equiv 0$ ,

# Examples

## Example

Let  $P(x_1, x_2, x_3) = x_1^2 x_2 - 2x_3$ . Pick any prime number  $p$  with  $p \equiv 3$  or  $p \equiv 5 \pmod{8}$ , so that 2 is not a quadratic residue modulo  $p$ . Then condition (1) is satisfied because  $z^3 - 2z \equiv 0$  iff  $z \equiv 0$ , and also condition (2) is easily verified.

# Examples

## Example

Let  $P(x_1, x_2, x_3) = x_1^2 x_2 - 2x_3$ . Pick any prime number  $p$  with  $p \equiv 3$  or  $p \equiv 5 \pmod{8}$ , so that 2 is not a quadratic residue modulo  $p$ . Then condition (1) is satisfied because  $z^3 - 2z \equiv 0$  iff  $z \equiv 0$ , and also condition (2) is easily verified. Since it has no constant solutions  $x_1 = x_2 = x_3$ , we can conclude that  $P(x_1, x_2, x_3)$  is not PR.

# Examples

## Example

Let  $P(x_1, x_2, x_3) = x_1^2 x_2 - 2x_3$ . Pick any prime number  $p$  with  $p \equiv 3$  or  $p \equiv 5 \pmod{8}$ , so that 2 is not a quadratic residue modulo  $p$ . Then condition (1) is satisfied because  $z^3 - 2z \equiv 0$  iff  $z \equiv 0$ , and also condition (2) is easily verified. Since it has no constant solutions  $x_1 = x_2 = x_3$ , we can conclude that  $P(x_1, x_2, x_3)$  is not PR.

Notice that, by Multiplicative Rado's Theorem, the seemingly similar equation  $x_1^2 x_2 = x_3$  is PR.

# Examples

## Example

Let  $P(x_1, x_2, x_3) = x_1^2 x_2 - 2x_3$ . Pick any prime number  $p$  with  $p \equiv 3$  or  $p \equiv 5 \pmod{8}$ , so that 2 is not a quadratic residue modulo  $p$ . Then condition (1) is satisfied because  $z^3 - 2z \equiv 0$  iff  $z \equiv 0$ , and also condition (2) is easily verified. Since it has no constant solutions  $x_1 = x_2 = x_3$ , we can conclude that  $P(x_1, x_2, x_3)$  is not PR.

Notice that, by Multiplicative Rado's Theorem, the seemingly similar equation  $x_1^2 x_2 = x_3$  is PR.

## Corollary

*Let  $P(\mathbf{x}) = \sum_{\alpha} c_{\alpha} \mathbf{x}^{\alpha} \in \mathbb{Z}[x_1, \dots, x_n]$  be an homogeneous polynomial. If for every nonempty  $J \subseteq \text{supp}(P)$  one has  $\sum_{\alpha \in J} c_{\alpha} \neq 0$ , then  $P(\mathbf{x})$  is not PR.*

# Necessary condition for sums of polynomials in one variable

## Theorem

*For every  $i = 1, \dots, n$  let  $P_i(x_i) = \sum_{s=1}^{d_i} c_{i,s} x_i^s$  be a polynomial of degree  $d_i$  in the variable  $x_i$  with no constant term.*



# Necessary condition for sums of polynomials in one variable

## Theorem

For every  $i = 1, \dots, n$  let  $P_i(x_i) = \sum_{s=1}^{d_i} c_{i,s} x_i^s$  be a polynomial of degree  $d_i$  in the variable  $x_i$  with no constant term. If the Diophantine equation

$$\sum_{i=1}^n P_i(x_i) = 0$$

is PR then the following “Rado’s condition” is satisfied:

# Necessary condition for sums of polynomials in one variable

## Theorem

For every  $i = 1, \dots, n$  let  $P_i(x_i) = \sum_{s=1}^{d_i} c_{i,s} x_i^s$  be a polynomial of degree  $d_i$  in the variable  $x_i$  with no constant term. If the Diophantine equation

$$\sum_{i=1}^n P_i(x_i) = 0$$

is PR then the following “Rado’s condition” is satisfied:

- “There exists a nonempty set  $J \subseteq \{1, \dots, n\}$  such that  $d_i = d_j$  for every  $i, j \in J$ , and  $\sum_{j \in J} c_{j,d_j} = 0$ .”

# Necessary condition for sums of polynomials in one variable

## Theorem

For every  $i = 1, \dots, n$  let  $P_i(x_i) = \sum_{s=1}^{d_i} c_{i,s} x_i^s$  be a polynomial of degree  $d_i$  in the variable  $x_i$  with no constant term. If the Diophantine equation

$$\sum_{i=1}^n P_i(x_i) = 0$$

is PR then the following “Rado’s condition” is satisfied:

- “There exists a nonempty set  $J \subseteq \{1, \dots, n\}$  such that  $d_i = d_j$  for every  $i, j \in J$ , and  $\sum_{j \in J} c_{j,d_j} = 0$ .”

Idea of the proof: by contradiction using  $p$ -expansions of hypernatural numbers; some refined nonstandard wizardry (overspilling principles, saturation) is used.

# Examples

# Examples

## Corollary

*A polynomial of the form  $\sum_{i=1}^n c_i x_i + P(y)$ , where  $P$  is a nonlinear polynomial with no constant term, is PR if and only if it is a Rado polynomial.*

# Examples

## Corollary

*A polynomial of the form  $\sum_{i=1}^n c_i x_i + P(y)$ , where  $P$  is a nonlinear polynomial with no constant term, is PR if and only if it is a Rado polynomial.*

## Example

The polynomial

$$P(x, y) = x^3 + 2x + y^3 - 2y$$

is not PR (even if it contains a partial sum of coefficients that equals zero).

# Examples

## Corollary

*A polynomial of the form  $\sum_{i=1}^n c_i x_i + P(y)$ , where  $P$  is a nonlinear polynomial with no constant term, is PR if and only if it is a Rado polynomial.*

## Example

The polynomial

$$P(x, y) = x^3 + 2x + y^3 - 2y$$

is not PR (even if it contains a partial sum of coefficients that equals zero).

## Example

The polynomials  $x^n + y^m = z^k$  are not PR for  $k \notin \{n, m\}$ .

# Open Problems/1



## Open Problems/1

**Open Problem 1.** Is  $x^2 + y^2 = z^2$  PR?

## Open Problems/1

**Open Problem 1.** Is  $x^2 + y^2 = z^2$  PR?

Recently, Heule, Kullmann and Marek have proven the PR of the Pythagorean equation for 2-colorings.

## Open Problems/1

**Open Problem 1.** Is  $x^2 + y^2 = z^2$  PR?

Recently, Heule, Kullmann and Marek have proven the PR of the Pythagorean equation for 2-colorings. Moreover, using number theoretical methods Chow, Lindqvist and Prendiville have proven the following:

## Open Problems/1

**Open Problem 1.** Is  $x^2 + y^2 = z^2$  PR?

Recently, Heule, Kullmann and Marek have proven the PR of the Pythagorean equation for 2-colorings. Moreover, using number theoretical methods Chow, Lindqvist and Prendiville have proven the following:

### Theorem

*For every  $k \in \mathbb{N}$  there exists  $s_0(k)$  such that for every  $s > s_0(k)$  and  $c_1, \dots, c_s \in \mathbb{Z} \setminus \{0\}$  the following equivalence holds:*

## Open Problems/1

**Open Problem 1.** Is  $x^2 + y^2 = z^2$  PR?

Recently, Heule, Kullmann and Marek have proven the PR of the Pythagorean equation for 2-colorings. Moreover, using number theoretical methods Chow, Lindqvist and Prendiville have proven the following:

### Theorem

*For every  $k \in \mathbb{N}$  there exists  $s_0(k)$  such that for every  $s > s_0(k)$  and  $c_1, \dots, c_s \in \mathbb{Z} \setminus \{0\}$  the following equivalence holds:*

- *The equation  $\sum_{i=1}^s c_i x_i^k$  is PR;*

## Open Problems/1

**Open Problem 1.** Is  $x^2 + y^2 = z^2$  PR?

Recently, Heule, Kullmann and Marek have proven the PR of the Pythagorean equation for 2-colorings. Moreover, using number theoretical methods Chow, Lindqvist and Prendiville have proven the following:

### Theorem

*For every  $k \in \mathbb{N}$  there exists  $s_0(k)$  such that for every  $s > s_0(k)$  and  $c_1, \dots, c_s \in \mathbb{Z} \setminus \{0\}$  the following equivalence holds:*

- *The equation  $\sum_{i=1}^s c_i x_i^k$  is PR;*
- *there exists a nonempty set  $J \subseteq \{1, \dots, s\}$  s.t.  $\sum_{j \in J} c_j = 0$ .*

## Open Problems/1

**Open Problem 1.** Is  $x^2 + y^2 = z^2$  PR?

Recently, Heule, Kullmann and Marek have proven the PR of the Pythagorean equation for 2-colorings. Moreover, using number theoretical methods Chow, Lindqvist and Prendiville have proven the following:

### Theorem

*For every  $k \in \mathbb{N}$  there exists  $s_0(k)$  such that for every  $s > s_0(k)$  and  $c_1, \dots, c_s \in \mathbb{Z} \setminus \{0\}$  the following equivalence holds:*

- *The equation  $\sum_{i=1}^s c_i x_i^k$  is PR;*
- *there exists a nonempty set  $J \subseteq \{1, \dots, s\}$  s.t.  $\sum_{j \in J} c_j = 0$ .*

$s_0(k)$  is of the order  $k \log k$ .

## Open Problems/1

**Open Problem 1.** Is  $x^2 + y^2 = z^2$  PR?

Recently, Heule, Kullmann and Marek have proven the PR of the Pythagorean equation for 2-colorings. Moreover, using number theoretical methods Chow, Lindqvist and Prendiville have proven the following:

### Theorem

*For every  $k \in \mathbb{N}$  there exists  $s_0(k)$  such that for every  $s > s_0(k)$  and  $c_1, \dots, c_s \in \mathbb{Z} \setminus \{0\}$  the following equivalence holds:*

- *The equation  $\sum_{i=1}^s c_i x_i^k$  is PR;*
- *there exists a nonempty set  $J \subseteq \{1, \dots, s\}$  s.t.  $\sum_{j \in J} c_j = 0$ .*

$s_0(k)$  is of the order  $k \log k$ .

**Open Problem 2.** Are there simple decidable conditions under which a given (non-homogeneous) Diophantine equation with no constant term is PR on  $\mathbb{N}$  if and only if it is PR on  $\mathbb{Z}$  if and only if it is PR on  $\mathbb{Q}$ ?



# Open Problems/2

## Open Problems/2

**Open Problem 3.** Are there simple “Rado-like” necessary and sufficient conditions under which a given Diophantine equation with no constant term is PR on sufficiently large finite fields  $\mathbb{F}_p$ ?

## Open Problems/2

**Open Problem 3.** Are there simple “Rado-like” necessary and sufficient conditions under which a given Diophantine equation with no constant term is PR on sufficiently large finite fields  $\mathbb{F}_p$ ?

**Open Problem 4** Is there a characterization of PR infinite systems of Diophantine equations in terms of  $u$ -equivalence? (Or, equivalently, by means of ultrafilters?)

# Thank You!

email: [lorenzo.luperi.baglini@univie.ac.at](mailto:lorenzo.luperi.baglini@univie.ac.at)